

LA-UR-19-21733

Approved for public release; distribution is unlimited.

Title: w18_plxa Viewgraphs

Author(s): Hsu, Scott C.
Langendorf, Samuel
Byvank, Tom
Stoltz, Peter H.

Intended for: Report

Issued: 2019-02-27

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

w18_plxa Viewgraphs

Scott Hsu, Samuel Langendorf, Tom Byvank, LANL
Peter Stoltz, Tech-X Corporation

2/28/2019



Operated by Triad National Security, LLC for the U.S. Department of Energy's NNSA

w18_plxa Viewgraphs

- Flash results from LA-UR-18-30344

Results presented in conference poster:

**T. Byvank, S. Langendorf, S. C. Hsu, P. Tzeferacos, Y. C. F. Thio,
Assessment of High- β Magnetized Target Formation and Plasma-Liner
Nonuniformities in Plasma-Jet-Driven Magneto-Inertial Fusion using the
FLASH Code, 60th Annual Meeting of the American Physical Society
Division of Plasma Physics (APS DPP 2018), Poster Session GP11,
Portland, OR, November 6, 2018.**

Background: Target Formation Quantities of Interest

Plasma Beta, $\beta = \Sigma_j n_j k T_j / (B^2 / 2\mu_0) \sim nT/B^2$

- Ratio of thermal pressure to magnetic pressure
- **GOAL: want $\beta > 1$** , thermally/flow-dominated
- High β limits MHD instabilities

(Ion) Magnetization, $\omega_i \tau_{ii} = \omega_i / \nu_{ii} \sim B T^{3/2} / n$

- Ratio of ion cyclotron frequency to ion-ion collision frequency
- **GOAL: want $\omega\tau > 1$** , magnetized (*despite high- β)
- High $\omega\tau$ reduces thermal transport \perp to B , enhances confinement time

Lambda Gun, $\lambda_{\text{gun}} = \mu_0 I / \Psi_{\text{pol}} = 2\pi r B_\theta / \int_0^R 2\pi r' B_z dr' \sim B_\theta / r_{\text{char}} B_z$

- In preliminary studies, we have $B_{\theta\text{MAX}} = 0.7$ T and we vary $B_{z0} = 0.004 - 0.4$ T


$\beta > 1$, $\omega\tau > 1$ is an interesting regime: thermally dominated and magnetized

Background: Lambda Gun Parameter

Spheromak

$$\lambda_{\text{gun}} > \lambda_{\text{crit}}$$

$$\beta < 1$$



Red: Plasma

Blue: Outside B

Green: Poloidal B


Yellow: Toroidal B

Wikihelper2134, 2017

Goal for Target

$$\infty > \lambda_{\text{gun}} > \lambda_{\text{crit}}$$

$$\beta > 1, \omega T > 1$$




Tangled magnetic field
(length scale $l \gg$ electron mean free path $\lambda_{\text{mfp,e}}$)
can further reduce electron transport
and increase confinement time

Hsu 2018 JFE, DOI: 10.1007/s10894-018-0168-z

Unmagnetized
Plasma Jet

$$\lambda_{\text{gun}} = \infty$$

$$\beta \gg 1, \omega T \ll 1$$




Optical
Emission
Image

Hsu 2012 PoP
DOI: 10.1063/1.4773320


Varying λ_{gun} will scan parameter space to create the $\beta > 1, \omega T > 1$ plasma

Simulation Setup: Target Formation

- 1) Plasma jets (H, 10 eV, $5 \cdot 10^{15} \text{ cm}^{-3}$) launched by $B_\theta(r,t) \sim \sin(r) \cdot \sin^2(t)$
 - 2) Jets advect $B_{\text{pol}}(r,z)$
 - 3) Merging jets form magnetized target
- FLASH Simulation: 2D azimuthally symmetric (r-z plane) of 2 colliding jets




Results: Target Formation– Time Evolution, Single Jet




$$\lambda_{\text{gun}} = 9 \text{ cm}^{-1}$$

Results: Target Formation– Time Evolution, Collision



$$\lambda_{\text{gun}} = 9 \text{ cm}^{-1}$$

Results: Target Formation– Varying λ_{gun}



Note: here, est. $\lambda_{\text{crit}} \sim 3/r_{\text{char}} = 0.6 \text{ cm}^{-1}$


Various λ_{gun} values can create the $\beta > 1$, $\omega\tau > 1$ plasma target

Simulation Setup: Liner Nonuniformities

Kim 2013 PoP
DOI: 10.1063/1.4789887


2D Discrete Jets

72 cm




15 cm jet length

2D Uniform Shell



9.6 cm shell thickness

3D Discrete Jets (60)







- Keeping total liner mass constant, 50 km/s (inward), 1.5 eV, 10^{17} cm^{-3}
- Initial target: 100 eV, $10^{18} \text{ cm}^{-3} \rightarrow 3.2 \cdot 10^4 \text{ bar}$




Cyan: Chamber vacuum (He), **Blue:** Liner (Ar), **Red:** Target (H)

Results: Liner Nonuniformities– Shocks

2D Discrete
Jets




2D Uniform
Shell




Merging of discrete jets creates shock structures. Question is: to what extent do these nonuniformities degrade the target compression?

Results: Liner Nonuniformities– Target Compression

2D Discrete Jets– Pressure




2D Uniform Shell– Pressure



Want small liner thickness so it acts like a piston to compress the target.

USim studies of 2D MHD target compressions


•Task 4.2.5: 2D PJMIF simulations



- We have the first simulations with axial magnetic field, consistent with (for example) embedding the target in a solenoidal field

USim studies of 2D MHD target compressions

•Task 4.2.5: 2D PJMIF simulations



- LANL supercomputing resources allowed us to run with sub-mm grid resolution with less than 30 min turnaround time